

PCT/EP2004/013162

18.11.2004



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 11 JAN 2005

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03078650.3

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

PCT/EP200 4 / 0 13 1 6 2

18.11.2004

Anmeldung Nr:
Application no.: 03078650.3
Demande no:

Anmeldetag:
Date of filing: 20.11.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

JANSSEN PHARMACEUTICA N.V.
Turnhoutseweg 30
2340 Beerse
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

7-phenylalkyl substituted 2-quinolinones and 2-quinoxalinones as poly(adp-ribose)
polymerase inhibitors

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07D241/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

7-PHENYLALKYL SUBSTITUTED 2-QUINOLINONES AND 2-QUINOXALINONES AS POLY(ADP-RIBOSE) POLYMERASE INHIBITORS

5 Field of the invention

The present invention relates to inhibitors of PARP and provides compounds and compositions containing the disclosed compounds. Moreover, the present invention provides methods of using the disclosed PARP inhibitors for instance as a medicine.

10 Background of the invention

The nuclear enzyme poly(ADP-ribose) polymerase-1 (PARP-1) is a member of the PARP enzyme family consisting of PARP-1 and several recently identified novel poly(ADP-ribosylating) enzymes. PARP is also referred to as poly(adenosine 5'-diphospho-ribose) polymerase or PARS (poly(ADP-ribose) synthetase).

15

PARP-1 is a major nuclear protein of 116 kDa consisting of three domains : the N-terminal DNA binding domain containing two zinc fingers, the automodification domain and the C-terminal catalytic domain. It is present in almost all eukaryotes. The enzyme synthesizes poly(ADP-ribose), a branched polymer that can consist of over 200 ADP-ribose units. The protein acceptors of poly(ADP-ribose) are directly or indirectly involved in maintaining DNA integrity. They include histones, topoisomerases, DNA and RNA polymerases, DNA ligases, and Ca^{2+} - and Mg^{2+} -dependent endonucleases. PARP protein is expressed at a high level in many tissues, most notably in the immune system, heart, brain and germ-line cells. Under normal physiological conditions, there is minimal PARP activity. However, DNA damage causes an immediate activation of PARP by up to 500-fold.

20

25

Among the many functions attributed to PARP, and especially PARP-1, is its major role in facilitating DNA repair by ADP-ribosylation and therefore co-ordinating a number of DNA repair proteins. As a result of PARP activation, NAD^+ levels significantly decline. Extensive PARP activation leads to severe depletion of NAD^+ in cells suffering from massive DNA damage. The short half-life of poly(ADP-ribose) results in a rapid turnover rate. Once poly(ADP-ribose) is formed, it is quickly degraded by the constitutively active poly(ADP-ribose) glycohydrolase (PARG), together with phosphodiesterase and (ADP-ribose) protein lyase. PARP and PARG form a cycle that converts a large amount of NAD^+ to ADP-ribose. In less than an hour, over-stimulation of PARP can cause a drop of NAD^+ and ATP to less than 20% of the normal level. Such a scenario is especially detrimental during ischaemia when

30

35

deprivation of oxygen has already drastically compromised cellular energy output. Subsequent free radical production during reperfusion is assumed to be a major cause of tissue damage. Part of the ATP drop, which is typical in many organs during ischaemia and reperfusion, could be linked to NAD^+ depletion due to poly(ADP-ribose) turnover. Thus, PARP or PARG inhibition is expected to preserve the cellular energy level thereby potentiating the survival of ischaemic tissues after insult.

Poly(ADP-ribose) synthesis is also involved in the induced expression of a number of genes essential for inflammatory response. PARP inhibitors suppress production of inducible nitric oxide synthase (iNOS) in macrophages, P-type selectin and intercellular adhesion molecule-1 (ICAM- 1) in endothelial cells. Such activity underlies the strong anti-inflammation effects exhibited by PARP inhibitors. PARP inhibition is able to reduce necrosis by preventing translocation and infiltration of neutrophils to the injured tissues.

PARP is activated by damaged DNA fragments and, once activated, catalyzes the attachment of up to 100 ADP-ribose units to a variety of nuclear proteins, including histones and PARP itself. During major cellular stresses the extensive activation of PARP can rapidly lead to cell damage or death through depletion of energy stores. As four molecules of ATP are consumed for every molecule of NAD^+ regenerated, NAD^+ is depleted by massive PARP activation, in the efforts to re-synthesize NAD^+ , ATP may also become depleted.

It has been reported that PARP activation plays a key role in both NMDA- and NO-induced neurotoxicity. This has been demonstrated in cortical cultures and in hippocampal slices wherein prevention of toxicity is directly correlated to PARP inhibition potency. The potential role of PARP inhibitors in treating neurodegenerative diseases and head trauma has thus been recognized even if the exact mechanism of action has not yet been elucidated.

Similarly, it has been demonstrated that single injections of PARP inhibitors have reduced the infarct size caused by ischemia and reperfusion of the heart or skeletal muscle in rabbits. In these studies, a single injection of 3-amino-benzamide (10 mg/kg), either one minute before occlusion or one minute before reperfusion, caused similar reductions in infarct size in the heart (32-42%) while 1,5- dihydroxyisoquinoline (1 mg/kg), another PARP inhibitor, reduced infarct size by a comparable degree (38-48%). These results make it reasonable to assume that PARP inhibitors could salvage previously ischaemic heart or reperfusion injury of skeletal muscle tissue.

PARP activation can also be used as a measure of damage following neurotoxic insults resulting from exposure to any of the following inducers like glutamate (via NMDA receptor stimulation), reactive oxygen intermediates, amyloid β -protein, N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) or its active metabolite N-methyl-4-phenylpyridine (MPP⁺), which participate in pathological conditions such as stroke, Alzheimer's disease and Parkinson's disease. Other studies have continued to explore the role of PARP activation in cerebellar granule cells in vitro and in MPTP neurotoxicity. Excessive neural exposure to glutamate, which serves as the predominate central nervous system neurotransmitter and acts upon the N-methyl D-aspartate (NMDA) receptors and other subtype receptors, most often occurs as a result of stroke or other neurodegenerative processes. Oxygen deprived neurons release glutamate in great quantities during ischaemic brain insult such as during a stroke or heart attack. This excess release of glutamate in turn causes over-stimulation (excitotoxicity) of N-methyl-D-aspartate (NMDA), AMPA, Kainate and MGR receptors, which open ion channels and permit uncontrolled ion flow (e.g., Ca²⁺ and Na⁺ into the cells and K⁺ out of the cells) leading to overstimulation of the neurons. The over-stimulated neurons secrete more glutamate, creating a feedback loop or domino effect which ultimately results in cell damage or death via the production of proteases, lipases and free radicals. Excessive activation of glutamate receptors has been implicated in various neurological diseases and conditions including epilepsy, stroke, Alzheimer's disease, Parkinson's disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's disease, schizophrenia, chronic pain, ischemia and neuronal loss following hypoxia, hypoglycemia, ischemia, trauma, and nervous insult. Glutamate exposure and stimulation has also been implicated as a basis for compulsive disorders, particularly drug dependence. Evidence includes findings in many animal species, as well as in cerebral cortical cultures treated with glutamate or NMDA, that glutamate receptor antagonists (i.e., compounds which block glutamate from binding to or activating its receptor) block neural damage following vascular stroke. Attempts to prevent excitotoxicity by blocking NMDA, AMPA, Kainate and MGR receptors have proven difficult because each receptor has multiple sites to which glutamate may bind and hence finding an effective mix of antagonists or universal antagonist to prevent binding of glutamate to all of the receptor and allow testing of this theory, has been difficult. Moreover, many of the compositions that are effective in blocking the receptors are also toxic to animals. As such, there is presently no known effective treatment for glutamate abnormalities. The stimulation of NMDA receptors by glutamate, for example, activates the enzyme neuronal nitric oxide synthase (nNOS), leading to the formation of nitric oxide (NO), which also mediates neurotoxicity. NMDA neurotoxicity may be prevented by

treatment with nitric oxide synthase (NOS) inhibitors or through targeted genetic disruption of nNOS in vitro.

5 Another use for PARP inhibitors is the treatment of peripheral nerve injuries, and the resultant pathological pain syndrome known as neuropathic pain, such as that induced by chronic constriction injury (CCI) of the common sciatic nerve and in which transsynaptic alteration of spinal cord dorsal horn characterized by hyperchromatosis of cytoplasm and nucleoplasm (so-called "dark" neurons) occurs.

10 Evidence also exists that PARP inhibitors are useful for treating inflammatory bowel disorders, such as colitis. Specifically, colitis was induced in rats by intraluminal administration of the hapten trinitrobenzene sulfonic acid in 50% ethanol. Treated rats received 3-aminobenzamide, a specific inhibitor of PARP activity. Inhibition of PARP activity reduced the inflammatory response and restored the morphology and the
15 energetic status of the distal colon.

Further evidence suggests that PARP inhibitors are useful for treating arthritis. Further, PARP inhibitors appear to be useful for treating diabetes. PARP inhibitors have been shown to be useful for treating endotoxic shock or septic shock.

20 PARP inhibitors have also been used to extend the lifespan and proliferative capacity of cells including treatment of diseases such as skin aging, Alzheimer's disease, atherosclerosis, osteoarthritis, osteoporosis, muscular dystrophy, degenerative diseases of skeletal muscle involving replicative senescence, age-related muscular degeneration,
25 immune senescence, AIDS, and other immune senescence disease; and to alter gene expression of senescent cells.

It is also known that PARP inhibitors, such as 3-amino benzamide, affect overall DNA repair in response, for example, to hydrogen peroxide or ionizing radiation.

30 The pivotal role of PARP in the repair of DNA strand breaks is well established, especially when caused directly by ionizing radiation or, indirectly after enzymatic repair of DNA lesions induced by methylating agents, topoisomerases I inhibitors and other chemotherapeutic agents as cisplatin and bleomycin. A variety of studies using
35 "knockout" mice, trans-dominant inhibition models (over-expression of the DNA-binding domain), antisense and small molecular weight inhibitors have demonstrated the role of PARP in repair and cell survival after induction of DNA damage. The

inhibition of PARP enzymatic activity should lead to an enhanced sensitivity of the tumor cells towards DNA damaging treatments.

5 PARP inhibitors have been reported to be effective in radiosensitizing (hypoxic) tumor cells and effective in preventing tumor cells from recovering from potentially lethal and sublethal damage of DNA after radiation therapy, presumably by their ability to prevent DNA strand break rejoining and by affecting several DNA damage signaling pathways.

10 PARP inhibitors have been used to treat cancer. In addition, U.S. Patent No.5,177,075 discusses several isoquinolines used for enhancing the lethal effects of ionizing radiation or chemotherapeutic agents on tumor cells. Weltin et al., "Effect of 6(5 - Phenanthridinone, an Inhibitor of Poly(ADP-ribose) Polymerase, on Cultured Tumor Cells", Oncol. Res., 6:9, 399-403 (1994), discusses the inhibition of PARP activity, reduced proliferation of tumor cells, and a marked synergistic effect when tumor cells
15 are co- treated with an alkylating drug.

A recent comprehensive review of the state of the art has been published by Li and Zhang in IDrugs 2001, 4(7): 804-812.


20 There continues to be a need for effective and potent PARP inhibitors, and more particularly PARP-1 inhibitors which produce minimal side effects. The present invention provides compounds, compositions for, and methods of, inhibiting PARP activity for treating cancer and/or preventing cellular, tissue and/or organ damage resulting from cell damage or death due to, for example, necrosis or apoptosis. The
25 compounds and compositions of the present invention are especially useful in enhancing the effectiveness of chemotherapy and radiotherapy where a primary effect of the treatment is that of causing DNA damage in the targeted cells.

Background prior art

30 EP 371564, published on June 6, 1990, discloses (1H-azol-1-ylmethyl) substituted quinoline, quinazoline or quinoxaline derivatives. The described compounds suppress the plasma elimination of retinoic acids. More in particular the compound 3-ethyl-7-[(1H-imidazol-1-ylphenylmethyl)-2(1H)-quinoxalinone (compound No
35 19 of the present application) was disclosed.

Description of the invention

This invention concerns compounds of formula (I)



5

the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

10 n is 0, 1 or 2;

X is N or CR^7 , wherein R^7 is hydrogen or taken together with R^1 may form a bivalent radical of formula $-CH=CH-CH=CH-$;

15 R^1 is C_{1-6} alkyl or thiophenyl;

R^2 is hydrogen, hydroxy, C_{1-6} alkyl, C_{3-6} alkynyl or taken together with R^3 may form $=O$;

R^3 is a radical selected from

- 20
- | | |
|----------------------|-----------|
| $-(CH_2)_s- NR^8R^9$ | (a-1), |
| $-O-H$ | (a-2), |
| $-O-R^{10}$ | (a-3), |
| $-S-R^{11}$ | (a-4), or |
| $-C\equiv N$ | (a-5), |

25 wherein

s is 0, 1, 2 or 3;

R^8 , R^{10} and R^{11} are each independently selected from $-CHO$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, piperidinyl, piperidinyl C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyl C_{1-6} alkyl, pyrrolyl C_{1-6} alkyl, aryl C_{1-6} alkylpiperidinyl, arylcarbonyl C_{1-6} alkyl, arylcarbonylpiperidinyl C_{1-6} alkyl, haloindazolylpiperidinyl C_{1-6} alkyl, or aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and

35 R^9 is hydrogen or C_{1-6} alkyl;



or R³ is a group of formula




wherein

t is 0, 1, 2 or 3;


5 Z is a heterocyclic ring system selected from



10



wherein R¹² is hydrogen, halo, C₁₋₆alkyl, aminocarbonyl, amino, hydroxy, aryl,



15

C₁₋₆alkylaminoC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkyloxyC₁₋₆alkylamino, arylC₁₋₆alkyl, di(phenylC₂₋₆alkenyl), piperidinyl, piperidinylC₁₋₆alkyl, C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkylC₁₋₆alkyl, aryloxy(hydroxy)C₁₋₆alkyl, haloindazolyl, arylC₁₋₆alkyl, arylC₂₋₆alkenyl, arylC₁₋₆alkylamino, morpholino, C₁₋₆alkylimidazolyl, or pyridinylC₁₋₆alkylamino;

20

R¹³ is hydrogen, piperidinyl or aryl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, halo, trihalomethyl, trihalomethoxy, C₁₋₆alkyl, C₁₋₆alkyloxy, amino, aminoC₁₋₆alkyl, di(C₁₋₆alkyl)amino, di(C₁₋₆alkyl)aminoC₁₋₆alkyloxy or C₁₋₆alkyloxycarbonyl, or C₁₋₆alkyl substituted with 1, 2 or 3 substituents independently selected from hydroxy, C₁₋₆alkyloxy, or aminoC₁₋₆alkyloxy; or

25

when R⁵ and R⁶ are on adjacent positions they may taken together form a bivalent radical of formula

- 5 -O-CH₂-O (d-1),
 -O-(CH₂)₂-O- (d-2),
 -CH=CH-CH=CH- (d-3), or
 -NH-C(O)-NR¹⁴=CH- (d-4),
 wherein R¹⁴ is C₁₋₆alkyl;

- 10 aryl is phenyl, phenyl substituted with halo, C₁₋₆alkyl or C₁₋₆alkyloxy;
 with the proviso that 3-ethyl-7-[(1*H*-imidazol-1-ylphenylmethyl]-2(1*H*)-quinoxalinone is not included.

- 15 Whenever the heterocyclic ring system Z contains a -CH₂-, -CH=, or -NH- moiety the substituents R¹² and R¹³ or the rest of the molecule can be attached to the carbon or nitrogen atom in which case one or both hydrogen atoms are replaced.

- 20 The compounds of formula (I) may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

A number of terms used in the foregoing definitions and hereinafter are explained hereunder. These terms are sometimes used as such or in composite terms.

- 25 As used in the foregoing definitions and hereinafter, halo is generic to fluoro, chloro, bromo and iodo; C₁₋₆alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, e.g. methyl, ethyl, propyl, butyl, pentyl, hexyl, 1-methylethyl, 2-methylpropyl, 2-methyl-butyl, 2-methylpentyl and the like; C₁₋₆alkanediyl defines bivalent straight and branched chained saturated
 30 hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,5-pentanediy, 1,6-hexanediyl and the branched isomers thereof such as, 2-methylpentanediyl, 3-methylpentanediyl, 2,2-dimethylbutanediyl, 2,3-dimethylbutanediyl and the like; trihaloC₁₋₆alkyl defines
 35 C₁₋₆alkyl containing three identical or different halo substituents for example trifluoromethyl; C₂₋₆alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 2 to 6 carbon atoms such as, for example, ethenyl, 2-propenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, and the like; C₃₋₆alkynyl defines straight and branch chained hydrocarbon radicals containing

one triple bond and having from 3 to 6 carbon atoms, such as, for example, 2-propynyl, 3-butyryl, 2-butyryl, 2-pentyryl, 3-pentyryl, 3-hexyryl, and the like; C₃₋₁₀cycloalkyl includes cyclic hydrocarbon groups having from 3 to 10 carbons, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cyclooctyl and the like.

The term "pharmaceutically acceptable salts" means pharmaceutically acceptable acid or base addition salts. The pharmaceutically acceptable acid or base addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and non-toxic base addition salt forms which the compounds of formula (I) are able to form. The compounds of formula (I) which have basic properties can be converted in their pharmaceutically acceptable acid addition salts by treating said base form with an appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic (*i.e.* butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

The compounds of formula (I) which have acidic properties may be converted in their pharmaceutically acceptable base addition salts by treating said acid form with a suitable organic or inorganic base. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-*D*-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The terms acid or base addition salt also comprise the hydrates and the solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The term stereochemically isomeric forms of compounds of formula (I), as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of formula (I) may possess. Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms which said compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of formula (I) both in pure form or in admixture with each other are

intended to be embraced within the scope of the present invention.

The *N*-oxide forms of the compounds of formula (I) are meant to comprise those compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide, particularly those *N*-oxides wherein one or more of the piperidine-,
5 piperazine or pyridazinyl-nitrogens are *N*-oxidized.

Whenever used hereinafter, the term "compounds of formula (I)" is meant to include also the *N*-oxide forms, the pharmaceutically acceptable acid or base addition salts and
10 all stereoisomeric forms.

The compounds described in EP 371564 suppress the plasma elimination of retinoic acids. 3-ethyl-7-[(1*H*-imidazol-1-ylphenyl)methyl]-2(1*H*)-quinoxalinone (compound No 19 of the present application) have been disclosed in EP 371564. Unexpectedly, it has
15 been found that the compounds of the present invention show PARP inhibitory activity.

A first group of interesting compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply:

- a) *n* is 0 or 1;
- 20 b) *X* is N or CR⁷, wherein R⁷ is hydrogen;
- c) R¹ is C₁₋₆alkyl;
- d) R² is hydrogen;
- e) R³ is a radical selected from (a-1) or (a-2) or is group of formula (b-1);
- f) *s* is 0, 1 or 2;
- 25 g) R⁸, R¹⁰ and R¹¹ are each independently selected from C₁₋₆alkyl or arylC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl;
- h) *t* is 0, 1 or 2;
- i) *Z* is a heterocyclic ring system selected from (c-1), (c-2), (c-3), (c-4), (c-5) or (c-11);
- j) R¹⁰ is hydrogen or C₁₋₆alkyloxyC₁₋₆alkylamino;
- 30 k) R¹¹ is hydrogen; and
- l) R⁴, R⁵ and R⁶ are each independently selected from hydrogen, halo or C₁₋₆alkyl.


A second group of interesting compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply:


- 35 a) *n* is 0 or 1;
- b) *X* is N;
- c) R¹ is C₁₋₆alkyl;
- d) R² is hydrogen;

- e) R^3 is a radical of formula (a-1) or is a group of formula (b-1);
 f) s is 0;
 g) R^8 is aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl;
 h) t is 0;
 5 i) Z is a heterocyclic ring system selected from (c-1) or (c-2);
 j) R^{12} is hydrogen or C_{1-6} alkyloxy C_{1-6} alkylamino;
 k) R^{13} is hydrogen; and
 k) R^4 , R^5 and R^6 are each independently selected from hydrogen or halo.
- 10 A group of preferred compounds consists of those compounds of formula (I) wherein n is 0 or 1; X is N or CR^7 , wherein R^7 is hydrogen; R^1 is C_{1-6} alkyl; R^2 is hydrogen; R^3 is a radical selected from (a-1) or (a-2) or is group of formula (b-1); s is 0, 1 or 2; R^8 is C_{1-6} alkyl or aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; t is 0, 1 or 2; Z is a heterocyclic ring system selected from (c-1), (c-2), (c-3), (c-4), (c-5) or (c-11); R^{12} is hydrogen or C_{1-6} alkyloxy C_{1-6} alkylamino; R^{13} is hydrogen; and R^4 , R^5 and R^6 are each independently selected from hydrogen, halo or C_{1-6} alkyl.
- 15

- A further group of preferred compounds consists of those compounds of formula (I) wherein n is 0 or 1; X is N; R^1 is C_{1-6} alkyl; R^2 is hydrogen; R^3 is a radical of formula (a-1) or is a group of formula (b-1); s is 0; R^8 is aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; t is 0; Z is a heterocyclic ring system selected from (c-1) or (c-2); R^{12} is hydrogen or C_{1-6} alkyloxy C_{1-6} alkylamino; R^{13} is hydrogen; and R^4 , R^5 and R^6 are each independently selected from hydrogen or halo.
- 20

- 25 The most preferred compounds are compound No 5, compound No 9, compound No 2 and compound No 1.







The compounds of formula (I) can be prepared according to the general methods described in EP 371564.

- 5 A number of such preparation methods will be described hereinafter in more detail. Other methods for obtaining final compounds of formula (I) are described in the examples.

- 10 Compounds of formula (I) wherein R² is hydrogen and R³ is -NR⁹-CHO wherein R⁹ is hydrogen or methyl, herein referred to as compounds of formula (I-b), can be prepared starting from compounds of formula (I), wherein R² taken together with R³ forms =O, herein referred to as compounds of formula (I-a), in the presence of formamide or methylformamide, here indicated as intermediates of formula (II), and formic acid.




- 20 Compounds of formula (I), wherein R³ is hydroxy, herein referred to as compounds of formula (I-c), can be prepared by converting the keton moiety of compounds of formula (I-a) into an hydroxy group, with an appropriate reductant, e.g., sodium borohydride in a suitable solvent, e.g. methanol and tetrahydrofuran.




Compounds of formula (I-a) can be prepared by converting compounds of formula (I-c), wherein R^2 is hydrogen, herein referred to as compounds of formula (I-c-1), in the presence of a suitable oxidant such as chromium trioxide and an acid such as sulfuric acid, in a suitable solvent such as 2-propanone.

5



Compounds of formula (I) wherein R^2 is hydrogen and R^3 is a radical of formula (c-1), herein referred to as a compound of formula (I-f), can be prepared by reacting compounds of formula (I) wherein R^2 is hydrogen and R^3 is a radical of formula (c-8), herein referred to as compounds of formula (I-d), with an amine of formula (III), wherein R^a is an appropriate radical, in the presence of a suitable solvent such as methanol and a suitable reagent such as sodium cyanoborohydride.


10




15

Intermediates of formula (IV), wherein W is an appropriate leaving group such as, for example, chloro, bromo, methanesulfonyloxy or benzenesulfonyloxy can be prepared from compounds of formula (I-c-1) by treating said compounds with a suitable reagent e.g. methanesulfonyloxy chloride or benzenesulfonyloxy chloride, or a halogenating reagent such as e.g. POCl_3 or SOCl_2 .

20



Compounds of formula (I), defined as compounds of formula (I) wherein R^b is as defined in R^8 and R^c is as defined in R^9 , or R^b and R^c taken together with the nitrogen to which they are attached, form an appropriate heterocyclic ring system as defined in Z, herein referred to as compounds of formula (I-h), can be prepared by reacting an intermediate of formula (IV) with an intermediate of formula (V). The reaction can be performed in a reaction-inert solvent such as dimethylformamide or acetonitrile, and optionally in the presence of a suitable base such as, for example, sodium carbonate, potassium carbonate or triethylamine.




The compounds of formula (I) may also be converted into each other via art-known reactions or functional group transformations. A number of such transformations are already described hereinabove. Other examples are hydrolysis of carboxylic esters to the corresponding carboxylic acid or alcohol; hydrolysis of amides to the corresponding carboxylic acids or amines; hydrolysis of nitriles to the corresponding amides; amino groups on imidazole or phenyl may be replaced by a hydrogen by art-known diazotation reactions and subsequent replacement of the diazo-group by hydrogen; alcohols may be converted into esters and ethers; primary amines may be converted into secondary or tertiary amines; double bonds may be hydrogenated to the corresponding single bond; an iodo radical on a phenyl group may be converted in to an ester group by carbon monoxide insertion in the presence of a suitable palladium catalyst.


25 Hence, compounds of formula (I), (I-a), (I-b), (I-c), (I-c-1), (I-d), (I-e), (I-f), (I-h), (I-i), (I-j) and (I-k) can optionally be the subject of one or more of the following conversions in any desired order:

- (i) converting a compound of formula (I) into a different compound of formula (I);
- (ii) converting a compound of formula (I) into the corresponding acceptable salt or N-oxide thereof;
- (iii) converting a pharmaceutically acceptable salt or N-oxide of a compound of formula (I) into the parent compound of formula (I);
- (iv) preparing a stereochemical isomeric form of a compound of formula (I) or a pharmaceutically acceptable salt or N-oxide thereof.

Intermediates of formula (VII), wherein R^d and R^e are appropriate radicals or taken together with the carbon to which they are attached, form an appropriate heterocyclic ring system as defined in Z, can be prepared by hydrolysing intermediates of formula (VI), wherein R^3 is a group of formula (b-1) or a radical of formula (a-1) wherein s is other than 0, herein referred to as R^B , according to art-known methods, such as stirring the intermediate (VI) in an aqueous acid solution in the presence of a reaction inert solvent, e.g. tetrahydrofuran. An appropriate acid is for instance hydrochloric acid.




Compounds of formula (I) wherein R^2 is hydrogen and R^B is as defined above, herein referred to as compounds of formula (I-k), can be prepared starting from intermediates of formula (VII), by a selective hydrogenation of said intermediate with an appropriate reducing agent such as, for example with a noble catalyst, such as platinum-on-charcoal, palladium-on-charcoal and the like and an appropriate reductant such as hydrogen in a suitable solvent such as methanol.




Compounds of formula (I) can be prepared by hydrolysing intermediates of formula (VIII), according to art-known methods, by submitting the intermediates of formula (VIII) to appropriate reagents, such as, tinchloride, acetic acid and hydrochloric acid, in the presence of a reaction inert solvent, e.g. tetrahydrofuran.

25




- 5 Compounds of formula (I) can be prepared starting from N-oxides of formula (IX) by converting the intermediates of formula (IX) in the presence of a suitable reagent such as sodium carbonate or acetic anhydride and when appropriate in a solvent such as dichloromethane.




- 10 The compounds of formula (I) wherein X is CH herein referred to as compounds of formula (I-j), may also be obtained by cyclizing an intermediate of formula (X). The cyclization reaction of intermediates of formula (X) may be conducted according to art-known cyclizing procedures. Preferably the reaction is carried out in the presence of a suitable Lewis Acid, e.g. aluminum chloride either neat or in a suitable solvent such as, for example, an aromatic hydrocarbon, e.g. benzene, chlorobenzene, methylbenzene and the like; halogenated hydrocarbons, e.g. trichloromethane, tetrachloromethane and the like; an ether, e.g. tetrahydrofuran, 1,4-dioxane and the like; or mixtures of such solvents. Somewhat elevated temperatures, preferably between 70°-100°C, and stirring may enhance the rate of the reaction.

20




The compounds of formula (I), wherein X is N, herein referred to as compounds of formula (I-i) may be obtained by condensing an appropriate ortho-benzenediamine of


- formula (XI) with an ester of formula (XII) wherein R^h is C_{1-6} alkyl. The condensation of the substituted ortho-diamine of formula (XI) and the ester of formula (XII) can be carried out in the presence of a carboxylic acid, e.g. acetic acid and the like, a mineral acid such as, for example hydrochloric acid, sulfuric acid, or a sulfonic acid such as, for example, methanesulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid and the like. Somewhat elevated temperatures may be appropriate to enhance the rate of the reaction and in some cases the reaction may even be carried out at the reflux temperature of the reaction mixture. The water which is liberated during the condensation may be removed from the mixture by azeotropical distillation, distillation and the like methods.




- Intermediates of formula (XI) can be prepared by a nitro to amine reduction reaction starting with an intermediate of formula (XIII) in the presence of a metal catalyst such as Raney Nickel and an appropriate reductant such as hydrogen in a suitable solvent such as methanol.



- Intermediates of formula (XIII) can be prepared by hydrolysing intermediates of formula (XIV), according to art-known methods, such as stirring the intermediate (XIV) in an aqueous acid solution in the presence of a reaction inert solvent, e.g. tetrahydrofuran,. An appropriate acid is for instance hydrochloric acid.




Intermediates of formula (X) can conveniently be prepared by reacting an aniline of formula (XV) with a halide of formula (XVI) in the presence of a base such as pyridine in a suitable solvent such as dichloromethane.



Intermediates of formula (VIII) wherein R^2 is hydrogen or hydroxy and when R^2 is hydrogen then R^3 is hydroxy herein referred to as intermediates of formula (VIII-a) can be prepared by treating an intermediate of formula (XVII), wherein W is halo, with an organolithium reagent such as, e.g. n-butyllithium in a reaction inert solvent, e.g. tetrahydrofuran, and subsequently reacting said intermediate with an intermediate of formula (XVIII) wherein R^1 is hydrogen or a radical as defined in R^3 .

10




The present invention also relates to a compound of formula (I) as defined above for use as a medicine.

The compounds of the present invention have PARP inhibiting properties as can be seen from the experimental part hereinunder.

20

The present invention contemplates the use of compounds in the preparation of a medicament for the treatment of any of the diseases and disorders in an animal described herein, wherein said compounds are compounds of formula (I)

25



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

5 n is 0, 1 or 2;

X is N or CR^7 , wherein R^7 is hydrogen or taken together with R^1 may form a bivalent radical of formula $-CH=CH-CH=CH-$;

10 R^1 is C_{1-6} alkyl or thiophenyl;

R^2 is hydrogen, hydroxy, C_{1-6} alkyl, C_{3-6} alkynyl or taken together with R^3 may form $=O$;

R^3 is a radical selected from

- 15 $-(CH_2)_s-NR^8R^9$ (a-1),
 $-O-H$ (a-2),
 $-O-R^{10}$ (a-3),
 $-S-R^{11}$ (a-4), or
 $-C\equiv N$ (a-5),


20 wherein

s is 0, 1, 2 or 3;

R^8 , R^{10} and R^{11} are each independently selected from $-CHO$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, piperidinyl, piperidinyl C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyl C_{1-6} alkyl, pyrrolyl C_{1-6} alkyl, aryl C_{1-6} alkylpiperidinyl, arylcarbonyl C_{1-6} alkyl, arylcarbonylpiperidinyl C_{1-6} alkyl, haloindazolylpiperidinyl C_{1-6} alkyl, or aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and

30 R^9 is hydrogen or C_{1-6} alkyl;


or R^3 is a group of formula



wherein


t is 0, 1, 2 or 3;

35 Z is a heterocyclic ring system selected from



5

wherein R^{12} is hydrogen, halo, C_{1-6} alkyl, aminocarbonyl, amino, hydroxy, aryl,



10


C_{1-6} alkylamino C_{1-6} alkyloxy, C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkylamino, aryl C_{1-6} alkyl, di(phenyl C_{2-6} alkenyl), piperidiny, piperidinyl C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl C_{1-6} alkyl, aryloxy(hydroxy) C_{1-6} alkyl, haloindazolyl, aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, aryl C_{1-6} alkylamino, morpholino, C_{1-6} alkylimidazolyl, or pyridinyl C_{1-6} alkylamino;
 R^{13} is hydrogen, piperidinyl or aryl;

15

R^4 , R^5 and R^6 are each independently selected from hydrogen, halo, trihalomethyl, trihalomethoxy, C_{1-6} alkyl, C_{1-6} alkyloxy, amino, amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl, or C_{1-6} alkyl substituted with 1, 2 or 3 substituents independently selected from hydroxy, C_{1-6} alkyloxy, or amino C_{1-6} alkyloxy; or
 when R^5 and R^6 are on adjacent positions they may taken together form a bivalent radical of formula

20

25



wherein R¹⁴ is C₁₋₆alkyl;

aryl is phenyl, phenyl substituted with halo, C₁₋₆alkyl or C₁₋₆alkyloxy.

- 5 In view of their PARP binding properties the compounds of the present invention may be used as reference compounds or tracer compounds in which case one of the atoms of the molecule may be replaced with, for instance, a radioactive isotope.

- 10 To prepare the pharmaceutical compositions of this invention, an effective amount of a particular compound, in base or acid addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by
- 15 parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders,
- 20 pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, to aid solubility for example, may be included. Injectable solutions, for
- 25 example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent,
- 30 optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect to the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. It is especially advantageous to
- 35 formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated

to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The compounds of the present invention can treat or prevent tissue damage resulting from cell damage or death due to necrosis or apoptosis; can ameliorate neural or cardiovascular tissue damage, including that following focal ischemia, myocardial infarction, and reperfusion injury; can treat various diseases and conditions caused or exacerbated by PARP activity; can extend or increase the lifespan or proliferative capacity of cells; can alter the gene expression of senescent cells; can radiosensitize and/or chemosensitize cells. Generally, inhibition of PARP activity spares the cells from energy loss, preventing, in the case of neural cells, irreversible depolarization of the neurons, and thus, provides neuroprotection.

For the foregoing reasons, the present invention further relates to a method of administering a therapeutically effective amount of the above-identified compounds in an amount sufficient to inhibit PARP activity, to treat or prevent tissue damage resulting from cell damage or death due to necrosis or apoptosis, to effect a neuronal activity not mediated by NMDA toxicity, to effect a neuronal activity mediated by NMDA toxicity, to treat neural tissue damage resulting from ischemia and reperfusion injury, neurological disorders and neurodegenerative diseases; to prevent or treat vascular stroke; to treat or prevent cardiovascular disorders; to treat other conditions and/or disorders such as age-related muscular degeneration, AIDS and other immune senescence diseases, inflammation, gout, arthritis, atherosclerosis, cachexia, cancer, degenerative diseases of skeletal muscle involving replicative senescence, diabetes, head trauma, inflammatory bowel disorders (such as colitis and Crohn's disease), muscular dystrophy, osteoarthritis, osteoporosis, chronic and/or acute pain (such as neuropathic pain), renal failure, retinal ischemia, septic shock (such as endotoxic shock), and skin aging, to extend the lifespan and proliferative capacity of cells; to alter gene expression of senescent cells; chemosensitize and/or radiosensitize (hypoxic) tumor cells. The present invention also relates to treating diseases and conditions in an animal which comprises administering to said animal a therapeutically effective amount of the above-identified compounds.

In particular, the present invention relates to a method of treating, preventing or inhibiting a neurological disorder in an animal, which comprises administering to said animal a therapeutically effective amount of the above-identified compounds. The

neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, traumatic brain injury, physical damage to the spinal cord, stroke associated with brain damage, focal ischemia, global ischemia, reperfusion injury, demyelinating disease and neurological disorder relating to
5 neurodegeneration.

The present invention also contemplates the use of compounds of formula (I) for inhibiting PARP activity, for treating, preventing or inhibiting tissue damage resulting from cell damage or death due to necrosis or apoptosis, for treating, preventing or
10 inhibiting a neurological disorder in an animal.

The term "preventing neurodegeneration" includes the ability to prevent neurodegeneration in patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for preventing further
15 neurodegeneration in patients who are already suffering from or have symptoms of a neurodegenerative disease.

The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes: (i) preventing a disease and/or
20 condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it; (ii) inhibiting the disease and/or condition, i.e., arresting its development; (iii) relieving the disease and/or condition, i.e., causing regression of the disease and/or condition.

25 The term "radiosensitizer", as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective amounts to increase the sensitivity of the cells to ionizing radiation and/or to promote the treatment of diseases which are treatable with ionizing radiation. Diseases which are treatable with ionizing radiation include neoplastic diseases, benign and malignant
30 tumors, and cancerous cells. Ionizing radiation treatment of other diseases not listed herein are also contemplated by the present invention.

The term "chemosensitizer", as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective
35 amounts to increase the sensitivity of cells to chemotherapy and/or promote the treatment of diseases which are treatable with chemotherapeutics. Diseases which are treatable with chemotherapy include neoplastic diseases, benign and malignant tumors

and cancerous cells. Chemotherapy treatment of other diseases not listed herein are also contemplated by the present invention.

5 The compounds, compositions and methods of the present invention are particularly useful for treating or preventing tissue damage resulting from cell death or damage due to necrosis or apoptosis.

The compounds of the present invention can be "anti-cancer agents", which term also encompasses "anti-tumor cell growth agents" and "anti-neoplastic agents". For example, the methods of the invention are useful for treating cancers and
10 chemosensitizing and/or radiosensitizing tumor cells in cancers such as ACTH-producing tumors, acute lymphocytic leukemia, acute nonlymphocytic leukemia, cancer of the adrenal cortex, bladder cancer, brain cancer, breast cancer, cervical cancer, chronic lymphocytic leukemia, chronic myelocytic leukemia, colorectal cancer, cutaneous T-cell lymphoma, endometrial cancer, esophageal cancer, Ewing's sarcoma
15 gallbladder cancer, hairy cell leukemia, head & neck cancer, Hodgkin's lymphoma, Kaposi's sarcoma, kidney cancer, liver cancer, lung cancer (small and/or non-small cell), malignant peritoneal effusion, malignant pleural effusion, melanoma, mesothelioma, multiple myeloma, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovarian cancer, ovary (germ cell) cancer, prostate cancer, pancreatic
20 cancer, penile cancer, retinoblastoma, skin cancer, soft tissue sarcoma, squamous cell carcinomas, stomach cancer, testicular cancer, thyroid cancer, trophoblastic neoplasms, uterine cancer, vaginal cancer, cancer of the vulva and Wilm's tumor.

Hence the compounds of the present invention can be used as "radiosensitizer" and/or
25 "chemosensitizer".

Radiosensitizers are known to increase the sensitivity of cancerous cells to the toxic effects of ionizing radiation. Several mechanisms for the mode of action of radiosensitizers have been suggested in the literature including: hypoxic cell
30 radiosensitizers (e.g., 2-nitroimidazole compounds, and benzotriazine dioxide compounds) mimicking oxygen or alternatively behave like bioreductive agents under hypoxia; non-hypoxic cell radiosensitizers (e.g., halogenated pyrimidines) can be analogs of DNA bases and preferentially incorporate into the DNA of cancer cells and thereby promote the radiation-induced breaking of DNA molecules and/or prevent the
35 normal DNA repair mechanisms; and various other potential mechanisms of action have been hypothesized for radiosensitizers in the treatment of disease.
Many cancer treatment protocols currently employ radiosensitizers in conjunction with radiation of x-rays. Examples of x-ray activated radiosensitizers include, but are not

limited to, the following: metronidazole, misonidazole, desmethylmisonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, EO9, RB 6145, nicotinamide, 5-bromodeoxyuridine (BUdR), 5-iododeoxyuridine (IUdR), bromodeoxycytidine, fluorodeoxyuridine (FudR), hydroxyurea, cisplatin, and therapeutically effective analogs and derivatives of the same.

5 Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, Photofrin, benzoporphyrin derivatives, tin etioporphyrin, pheorbide-a, bacteriochlorophyll-a, 10 naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

Radiosensitizers may be administered in conjunction with a therapeutically effective amount of one or more other compounds, including but not limited to: compounds which promote the incorporation of radiosensitizers to the target cells; compounds 15 which control the flow of therapeutics, nutrients, and/or oxygen to the target cells; chemotherapeutic agents which act on the tumor with or without additional radiation; or other therapeutically effective compounds for treating cancer or other disease. Examples of additional therapeutic agents that may be used in conjunction with radiosensitizers include, but are not limited to: 5-fluorouracil, leucovorin, 5'-amino- 20 5'deoxythymidine, oxygen, carbogen, red cell transfusions, perfluorocarbons (e.g., Fluosol 10 DA), 2,3-DPG, BW12C, calcium channel blockers, pentoxifylline, antiangiogenesis compounds, hydralazine, and LBSO. Examples of chemotherapeutic agents that may be used in conjunction with radiosensitizers include, but are not limited to: adriamycin, camptothecin, carboplatin, cisplatin, daunorubicin, docetaxel, 25 doxorubicin, interferon (alpha, beta, gamma), interleukin 2, irinotecan, paclitaxel, topotecan, and therapeutically effective analogs and derivatives of the same.

Chemosensitizers may be administered in conjunction with a therapeutically effective amount of one or more other compounds, including but not limited to : compounds 30 which promote the incorporation of chemosensitizers to the target cells; compounds which control the flow of therapeutics, nutrients, and/or oxygen to the target cells; chemotherapeutic agents which act on the tumor or other therapeutically effective compounds for treating cancer or other disease. Examples of additional therapeutical agents that may be used in conjunction with chemosensitizers include, but are not 35 limited to : methylating agents, topoisomerase I inhibitors and other chemotherapeutic agents such as cisplatin and bleomycin.

The compounds of formula (I) can also be used to detect or identify the PARP, and

more in particular the PARP-1 receptor. For that purpose the compounds of formula (I) can be labeled. Said label can be selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label fluorescent group or a chemiluminiscent group.

- 5 Those skilled in the art could easily determine the effective amount from the test results presented hereinafter. In general it is contemplated that an effective amount would be from 0.001 mg/kg to 100 mg/kg body weight, and in particular from 0.005 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses
10 may be formulated as unit dosage forms, for example, containing 0.05 to 500 mg, and in particular 0.1 mg to 200 mg of active ingredient per unit dosage form.

The following examples illustrate the present invention.


15 Experimental part

Hereinafter, "BuLi" is defines as butyl-lithium, "MeOH" is defined as methanol, "DIPE" is defined as diisopropyl ether, "DMF" is defined as *N,N*-dimethylformamide, "DME" is defined as 1,2-dimethoxyethane, "DCM" is defined as dichloromethane, "EtOAc" is defined as ethyl acetate, "THF" is defined as tetrahydrofuran, "MEK" is
20 defined as methyl ethyl keton.

A. Preparation of the intermediate compounds


Example A1

a) Preparation of intermediate 1



- 25 Nitric acid (fuming) (26.7ml) was added dropwise at room temperature to a solution of *N*-[4-(2-oxo-2-phenylethyl)phenyl]- acetamide (0.2128 mol) in acetic acid, anhydride (1100ml) while the temperature was kept below 30°C. The mixture was stirred for 1 hour, poured out into ice water and neutralized with a concentrated NH_4OH solution. The precipitate was filtered off, washed with water and with diethyl ether and dried, yielding 40g (63%) of intermediate 1.


b) Preparation of intermediate 2



Sodium hydroborate (0.1056 mol) was added portionwise at 10°C under N₂ flow to a solution of intermediate 1 (0.096 mol) in methanol (350ml). The mixture was stirred at 10°C for 30 min, poured out into ice water and extracted with DCM. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness,

5 yielding 22g (76%) of intermediate 2.


c) Preparation of intermediate 3



Methylsulfonyl chloride (0.076 mol) was added dropwise at 0°C under N₂ flow to a suspension of intermediate 2 (0.038 mol) and triethylamine (0.076 mol) in DCM (100ml). The mixture was stirred for 12 hours. The solvent was evaporated (without heating). The product was used without further purification, yielding (quant.) of

10 intermediate 3.


d) Preparation of intermediate 4



A mixture of intermediate 3 (0.038 mol), 1H-imidazole (0.076 mol) and potassium carbonate (0.076 mol) in acetonitrile (300ml) was stirred and refluxed for 15 hours, then cooled to room temperature, poured out into ice water and extracted with DCM.


15 The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH/NH₄OH 95/5/0.1). The pure fractions were collected and the solvent was evaporated, yielding 4.5g (34%) of intermediate 4.

e) Preparation of intermediate 5



20 A mixture of intermediate 4 (0.0128 mol) in sodium hydroxide (110ml) and ethanol (30ml) was stirred at room temperature for 3 hours, poured out on ice and extracted with DCM. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness, yielding 2.75g (70%) of intermediate 5.


f) Preparation of intermediate 6



25 A mixture of intermediate 5 (0.0089 mol) in methanol (150ml) was hydrogenated under a 3 bar pressure for 45 min with Raney Nickel (3g) as a catalyst. After uptake of H₂ (3 equiv), the catalyst was filtered through celite and the filtrate was evaporated till dryness, yielding 2.59g (100%) of intermediate 6.


Example A2

a) Preparation of intermediate 7




- 5 A mixture of 3,4-benzophenonediamine (0.1743 mol) and ethyl-2-oxobutanoate (0.3486 mol) in ethanol (820ml) was stirred and refluxed for 5 hours and then cooled. The solvent was evaporated. The residue was taken up in an aqueous saturated NaHCO_3 solution. The mixture was extracted with DCM. The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated till dryness. The residue (70.3g) was purified by column chromatography over silica gel (eluent: DCM/2-propanol 98/2; 20-45 5m). The pure fractions were collected and the solvent was evaporated, yielding 15.5g of intermediate 7. Part of it was crystallized from diethyl ether and petroleum ether. The precipitate was filtered off and dried, yielding 0.85g of intermediate 7, melting point 197°C .

b) Preparation of intermediate 8




- 15 Sodium hydroborate (0.0719 mol) was added at 0°C under N_2 flow to a suspension of intermediate 7 ((0.0719 mol in methanol (300ml) and THF (100ml)). The mixture was stirred for 30 min and poured out into ice water. The precipitate was filtered off, washed with water and dried. A part (0.7g) of this fraction (8.33g) was crystallized from methanol and water. The precipitate was filtered off and dried, yielding 0.59g of intermediate 8, melting point 202°C .

c) Preparation of intermediate 9



- 20 Methanesulfonyl chloride (0.009421 mol) was added at 0°C under N_2 flow to a suspension of intermediate 8 (0.0043 mol) and triethylamine (0.0108 mol) in DCM (10ml) and THF (10ml). The mixture was stirred at room temperature for 5 hours. The solvent was evaporated cold. The product was used without further purification, yielding (100%) of intermediate 9.

d) Preparation of intermediate 10




- 25 A mixture of *N*-(2-methoxyethyl)-1-(phenylmethyl)-4-piperidinamine (0.0402 mol) in ethanol (100ml) was hydrogenated at 40°C for 2 hours and then at room temperature under a 3 bar pressure for 3 hours with Pd/C 10% (1g) as a catalyst. After uptake of H_2 (1 equiv), the catalyst was filtered through celite, washed with ethanol and the filtrate

was evaporated. The product was used without further purification, yielding 6.5g (99%) of intermediate 10.


Example A3

a) Preparation of intermediate 11




- 5 A mixture of 6-bromo-2-chloro-3-methyl- quinoline (0.0697 mol) and NaOCH₃ 30% (0.3483 mol) in methanol (90ml) was stirred and refluxed for 15 hours. The mixture was cooled, poured out into ice water and extracted with DCM. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The product was used without further purification, yielding 12.2g (69%) of intermediate 11.

b) Preparation of intermediate 12




- 10 n-BuLi (0.0624 mol) was added dropwise at -60°C under N₂ flow to a solution of intermediate 11 (0.048 mol) in THF (100ml) and the mixture was stirred at -60°C for 1 hour. A solution of 2-benzoyldimethylamine (0.0576 mol) in THF (100ml) was added dropwise. The mixture was allowed to warm to -20°C while stirring and then stirred for 2 hours. The mixture was poured out into an aqueous NH₄Cl solution and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The product was used without further purification, yielding 17.68g (quant.) of intermediate 12.

c) Preparation of intermediate 13



- 20 A mixture of intermediate 12 (0.0496 mol) in HCl 3N (261ml) and THF (133ml) was stirred and refluxed for 8 hours. The mixture was cooled, poured out on ice, basified with a concentrated NH₄OH solution and extracted with DCM. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The residue was purified by column chromatography over silica gel (15-40 μm) (eluent: DCM/CH₃OH/NH₄OH 95/5/1). The pure fractions were collected and the solvent was evaporated, yielding 6g (36%) of intermediate 13.

d) Preparation of intermediate 14




- 25 A mixture of intermediate 13 (0.0178 mol) in HCl 3N (90ml) and THF (47ml) was stirred and refluxed for 2 days. The mixture was cooled, poured out on ice, basified

with a concentrated NH_4OH solution and extracted with DCM. The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated till dryness. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried, yielding 1.5g of intermediate 14.

5

Example A4


a) Preparation of intermediate 15



A mixture of (4-amino-3-nitrophenyl)(4-chlorophenyl)-methanone (0.0686 mol) in DCM (200 ml) and acetyl chloride (20 ml) was stirred for 12 hours at room temperature and then the solvent was evaporated dry. The residue was taken up in diethyl ether (50 ml), then the desired product was filtered off and dried, yielding 21.6 g (99 %) of intermediate 15, melting point 138°C .

10


b) Preparation of intermediate 16



A mixture of intermediate 15 (0.066 mol) in methanol (200 ml) was stirred at 0°C and a solution of sodium hydroborate (0.066 mol) in water was added dropwise, then the reaction mixture was stirred for 1 hour at room temperature and the solvent was evaporated. The residue was extracted with DCM/ $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ and the extract was dried (MgSO_4). Finally the solvent was evaporated and the desired product was collected, yielding 20.4 g (97 %) of intermediate 16, melting point 198°C .

15


c) Preparation of intermediate 17



In a 3-neck reaction flask (500 ml), equipped with an addition funnel and thermometer, a mixture of intermediate 16 (0.062 mol) and triethylamine (0.125 mol) in DCM (200 ml) was cooled to 0°C and methanesulfonyl chloride (0.125 mol) was added dropwise keeping the temperature at $0-5^\circ\text{C}$, then the reaction mixture was stirred for 4 hours at room temperature and poured out into water (1000 ml). The organic layer was separated, dried (MgSO_4), filtered off and the solvent was evaporated, yielding 18 g (oil, 85 %) of intermediate 17.


20

d) Preparation of intermediate 18




- 5 A mixture of intermediate 17 (0.088 mol), piperidine (0.446 mol) and potassium carbonate (0.442 mol) in acetonitrile (250 ml) with a small amount of KI was stirred at 40°C for 12 hours and the solvent was evaporated (vac.). The residue was taken up in water and the mixture was extracted with DCM. The organic layer was dried (MgSO₄) and the solvent was evaporated dry. The residue was purified by liquid chromatography over silica gel (eluent: DCM). The product fractions were collected and the solvent was evaporated, yielding 16 g (47 %, oil) of intermediate 18.

e) Preparation of intermediate 19



- 10 A mixture of intermediate 18 (0.0413 mol) in sodium hydroxide (1.5N) (160 ml) and THF/methanol (10 ml) was stirred for 48 hours at room temperature, then the solution was neutralised to pH: 7, extracted with EtOAc and washed with water. The organic layer was dried (MgSO₄) and the solvent was evaporated dry. The oily residue (12.5 g) was purified by high-performance liquid chromatography over silica gel (eluent: DCM/CH₃OH 99/1). The product fractions were collected and the solvent was evaporated, yielding 10 g (71 %) of intermediate 19, melting point 124°C.

h) Preparation of intermediate 20




- 15 A mixture of intermediate 19 (0.0289 mol) in methanol (200 ml) was hydrogenated for 2 hours with Raney Nickel (10 g) as a catalyst. After uptake of H₂ (3 equiv.), the solution was filtered over a celite path, yielding 9.1 g of intermediate 20 (used as such in the next reaction step without further purification).

20 B. Preparation of the final compounds

Example B1

Preparation of compound 1




A mixture of intermediate 6 (0.0089 mol) and ethyl 2-oxobutanoate (0.0178 mol) in methanol (50ml) was stirred and refluxed for 15 hours, then cooled to room temperature, poured out into ice water and extracted with DCM. The organic layer was

separated, dried (MgSO_4), filtered and the solvent was evaporated till dryness. The residue (3.1g) was purified by column chromatography over silica gel (15-40 μm) (eluent: toluene/2-propanol/ NH_4OH 85/15/0.8). The desired fractions were collected and the solvent was evaporated. The residue was purified by column chromatography over silica gel (15-40 μm)(eluent: toluene/2-propanol/ NH_4OH 85/15/0.8). The pure fractions were collected and their solvents were evaporated. The residue (0.3g) was crystallized from 2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 0.3g (10%) of compound 1, melting point 166°C.

10 Example B2

Preparation of compound 2




. $\text{C}_2\text{H}_2\text{O}_4$ (2:5)

A mixture of intermediate 9 (0.0043 mol), intermediate 10 (0.0052 mol) and potassium carbonate (0.0129 mol) in acetonitrile (15ml) was stirred at 80°C for 15 hours, then cooled to room temperature, poured out into ice water and extracted with EtOAc. The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated till dryness. The residue (2:2g) was purified by column chromatography over silica gel (15-40 μm) (eluent: DCM/ CH_3OH / NH_4OH 95/5/0.2). The pure fractions were collected and the solvent was evaporated. The residue (0.27g) was dissolved in 2-propanone and converted into the ethanedioic acid salt (2:5). The precipitate was filtered off and dried, yielding 0.25g of compound 2, melting point 98°C.

20

Example B3

Preparation of compound 3




Intermediate 14 (0.0107 mol) in methanol (60ml) was hydrogenated with Pd/C 10% (0.36g) as a catalyst for 16h under a 3 bar pressure. After uptake of H_2 (1eq), the catalyst was filtered through celite and the filtrate was evaporated till dryness. The residue (3.58g) was purified by column chromatography over silica gel (15-40 μm) (eluent : DCM/ CH_3OH / NH_4OH 93/7/0.5). The pure fractions were collected and the solvent was evaporated. The residue (1.45g) was crystallized from methyl ethyl keton. The precipitate was filtered off, washed with diethylether and dried, yielding 0.82g (30%) of compound 3.

30

Example B4


Preparation of compound 4



- Sodium hydroborate (0.0719 mol) was added at 0°C under N₂ flow to a suspension of intermediate 7 (0.0719 mol) in methanol (300ml) and THF (100ml). The mixture was stirred for 30 min and poured out into ice water. The precipitate was filtered off, washed with water and dried. A part (0.7g) of this fraction (8.33g) was crystallized from methanol and water. The precipitate was filtered off and dried, yielding 0.59g of compound 4, melting point 202°C.

Example B5















Preparation of compound 5






- A solution of 2-oxo-butanoic acid (0.0294 mol) in acetic acid (30 ml) was added to a solution of intermediate 20 (0.0288 mol) in water (70 ml) at 0°C and the reaction mixture was stirred for 2 hours at 0°C. The resulting solution was poured out into ice-water, neutralised with sodium hydroxide (3N) and extracted with DCM. The organic layer was dried (MgSO₄) and the solvent was evaporated dry. The residue was purified by liquid chromatography over silica gel (eluent: toluene/2-propanol/NH₄OH 90/10/0.1). Two product fractions were collected and the solvent was evaporated. The upper fraction after purification was crystallised from diethylether/DCM/MEK and the desired product was collected, yielding 1.15 g of compound 5.
- Table F-1 lists the compounds that were prepared according to one of the above Examples. The following abbreviations were used in the tables: Co.No. stands for Compound Number, Ex. [Bn°] referred to the same method as described in the Bn° examples.

Table 1:

Co. No. 1; Ex. [B1]; mp. 166°C.	C ₂ H ₂ O ₄ (2:5); Co. No. 2; Ex. [B2]; mp. 98°C

	
Co. No. 3; Ex. [B3]	Co. No. 4; Ex. [B4]; mp. 202°C
	
Co. No. 5; Ex. [B5]	Co. No. 6; Ex. [B1]; mp. 182°C
	
Co. No. 7; Ex. [B1]; mp. 176°C	Co. No. 8; Ex. [B1]; mp. 210.4°C
	
C ₂ H ₂ O ₄ (1:2); Co. No. 9; Ex. [B2]; mp. 118°C	Co. No. 10; Ex. [B4]; mp. 254°C
	
Co. No. 11; Ex. [B4]; mp. 202°C	Co. No. 12; Ex. [B5]; mp. 174.6°C
	
Co. No. 13; Ex. [B5]; mp. 203.1°C	Co. No. 14; Ex. [B5]; mp. 230°C
	
Co. No. 15; Ex. [B5]; mp. 184°C	Co. No. 16; Ex. [B5]

	
<p>Co. No. 17; Ex. [B5]; mp. 264.4°C</p>	<p>Co. No. 18; Ex. [B5]; mp. 118°C</p>
	
<p>EP0371564; Co. No. 19</p>	

Pharmacological example

In vitro Scintillation Proximity Assay (SPA) for PARP-1 inhibitory activity

Compounds of the present invention were tested in an *in vitro* assay based on SPA

5 technology (proprietary to Amersham Pharmacia Biotech).

In principle, the assay relies upon the well established SPA technology for the detection of poly(ADP-ribosylation) of biotinylated target proteins, i.e histones. This ribosylation is induced using nicked DNA activated PARP-1 enzyme and [³H]-nicotinamide adenine dinucleotide ([³H]-NAD⁺) as ADP-ribosyl donor.

10

As inducer of PARP-1 enzyme activity, nicked DNA was prepared. For this, 25 mg of DNA (supplier: Sigma) was dissolved in 25 ml DNase buffer (10 mM Tris-HCl, pH 7.4; 0.5 mg/ml Bovine Serum Albumine (BSA); 5 mM MgCl₂·6H₂O and 1 mM KCl) to which 50 µl DNase solution (1mg/ml in 0.15 M NaCl) was added. After an incubation

15

of 90 min. at 37 °C, the reaction was terminated by adding 1.45 g NaCl, followed by a further incubation at 58 °C for 15 min. The reaction mixture was cooled on ice and dialysed at 4 °C for respectively 1.5 and 2 hours against 1.5 l of 0.2 M KCl, and twice

20

against 1.5 l of 0.01 M KCl for 1.5 and 2 h respectively. The mixture was aliquoted and stored at -20 °C. Histones (1 mg/ml, type II-A, supplier: Sigma) were biotinylated using the biotinylation kit of Amersham and stored aliquoted at -20 °C. A stock solution of 100 mg/ml SPA poly(vinyl toluene) (PVT) beads (supplier: Amersham) was made in PBS. A stock solution of [³H]-NAD⁺ was made by adding 120 µl of [³H]-

25

NAD⁺ (0.1 mCi/ml, supplier: NEN) to 6 ml incubation buffer (50 mM Tris/HCl, pH 8; 0.2 mM DTT; 4 mM MgCl₂). A solution of 4 mM NAD⁺ (supplier: Roche) was made in incubation buffer (from a 100 mM stock solution in water stored at -20 °C). The PARP-1 enzyme was produced using art known techniques, i.e. cloning and expression of the protein starting from human liver cDNA. Information concerning the used protein sequence of the PARP-1 enzyme including literature references can be found in the Swiss-Prot database under primary accession number P09874. Biotinylated histones

30

and PVT-SPA beads were mixed and pre-incubated for 30 min. at room temperature. PARP-1 enzyme (concentration was lot dependent) was mixed with the nicked DNA and the mixture was pre-incubated for 30 min. at 4 °C. Equal parts of this

35

histones/PVT-SPA beads solution and PARP-1 enzyme/DNA solution were mixed and 75 µl of this mixture together with 1 µl of compound in DMSO and 25 µl of [³H]-NAD⁺ was added per well into a 96-well microtiterplate. The final concentrations in the incubation mixture were 2 µg/ml for the biotinylated histones, 2 mg/ml for the PVT-SPA beads, 2 µg/ml for the nicked DNA and between 5 - 10 µg/ml for the PARP-1 enzyme. After incubation of the mixture for 15 min. at room temperature, the reaction

was terminated by adding 100 μ l of 4 mM NAD^+ in incubation buffer (final concentration 2 mM) and plates were mixed.

The beads were allowed to sediment for at least 15 min. and plates transferred to a TopCountNXTTM (Packard) for scintillation counting, values were expressed as counts per minute (cpm). For each experiment, controls (containing PARP-1 enzyme and DMSO without compound), a blank incubation (containing DMSO but no PARP-1 enzyme or compound) and samples (containing PARP-1 enzyme and compound dissolved in DMSO) were run in parallel. All compounds tested were dissolved and eventually further diluted in DMSO. In first instance, compounds were tested at a concentration of 10^{-5} M. When the compounds showed activity at 10^{-5} M, a dose-response curve was made wherein the compounds were tested at concentrations between 10^{-5} M and 10^{-8} M. In each test, the blank value was subtracted from both the control and the sample values. The control sample represented maximal PARP-1 enzyme activity. For each sample, the amount of cpm was expressed as a percentage of the mean cpm value of the controls. When appropriate, IC_{50} -values (concentration of the drug, needed to reduce the PARP-1 enzyme activity to 50% of the control) were computed using linear interpolation between the experimental points just above and below the 50 % level. Herein the effects of test compounds are expressed as pIC_{50} (the negative log value of the IC_{50} -value). As a reference compound, 4-amino-1,8-naphthalimide was included to validate the SPA assay. The tested compounds showed inhibitory activity at the initial test concentration of 10^{-5} M (see Tabel-2).

In vitro filtration assay for PARP-1 inhibitory activity

Compounds of the present invention were tested in an *in vitro* filtration assay assessing PARP-1 activity (triggered in the presence of nicked DNA) by means of its histone poly (ADP-ribosyl)ation activity using [^{32}P]-NAD as ADP-ribosyl donor. The radioactive ribosylated histones were precipitated by trichloroacetic acid (TCA) in 96-well filterplates and the incorporated [^{32}P] measured using a scintillation counter


A mixture of histones (stock solution: 5 mg/ml in H_2O), NAD^+ (stock solution: 100 mM in H_2O), and [^{32}P]- NAD^+ in incubation buffer (50 mM Tris/HCl, pH 8; 0.2 mM DTT; 4 mM MgCl_2) was made. A mixture of the PARP-1 enzyme (5 – 10 μ g/ml) and nicked DNA was also made. The nicked DNA was prepared as described in the *in vitro* SPA for PARP-1 inhibitory activity. Seventy-five μ l of the PARP-1 enzyme/DNA mixture together with 1 μ l of compound in DMSO and 25 μ l of histones- NAD^+ /[^{32}P]- NAD^+ mixture was added per well of a 96-well filterplate (0.45 μ m, supplier Millipore). The final concentrations in the incubation mixture were 2 μ g/ml for the

histones, 0.1 mM for the NAD^+ , 200 μM (0.5 μC) for the $[\text{}^{32}\text{P}]\text{-NAD}^+$ and 2 $\mu\text{g/ml}$ for the nicked DNA. Plates were incubated for 15 min. at room temperature and the reaction was terminated by the addition of 10 μl ice cold 100% TCA followed by the addition of 10 μl ice-cold BSA solution (1 % in H_2O). The protein fraction was allowed to precipitate for 10 min. at 4 $^\circ\text{C}$ and plates were vacuum filtered. The plates were subsequently washed with, for each well, 1 ml of 10 % ice cold TCA, 1 ml of 5 % ice cold TCA and 1 ml of 5 % TCA at room temperature. Finally 100 μl of scintillation solution (Microscint 40, Packard) was added to each well and the plates were transferred to a TopCountNXTTM (supplier: Packard) for scintillation counting and values were expressed as counts per minute (cpm). For each experiment, controls (containing PARP-1 enzyme and DMSO without compound), a blank incubation (containing DMSO but no PARP-1 enzyme or compound) and samples (containing PARP-1 enzyme and compound dissolved in DMSO) were run in parallel. All compounds tested were dissolved and eventually further diluted in DMSO. In first instance, compounds were tested at a concentration of 10^{-5}M . When the compounds showed activity at 10^{-5}M , a dose-response curve was made wherein the compounds were tested at concentrations between 10^{-5}M and 10^{-8}M . In each test, the blank value was subtracted from both the control and the sample values. The control sample represented maximal PARP-1 enzyme activity. For each sample, the amount of cpm was expressed as a percentage of the mean cpm value of the controls. When appropriate, IC_{50} -values (concentration of the drug, needed to reduce the PARP-1 enzyme activity to 50% of the control) were computed using linear interpolation between the experimental points just above and below the 50 % level. Herein the effects of test compounds are expressed as pIC_{50} (the negative log value of the IC_{50} -value). As a reference compound, 4-amino-1,8-naphthalimide was included to validate the filtration assay. The tested compounds showed inhibitory activity at the initial test concentration of 10^{-5}M (see Tabel-2).

Tabel-2


Co.No.	In vitro SPA	In vitro filtration assay
	pIC50	pIC50
12	6.138	5.444
13	6.144	5.306
14	5.797	5.573
5	6.628	6.159
15	6.483	5.195
16	6.239	5.344
17	6.476	
9	6.506	
18	6	5.384
19	6.636	6.211
2	8.125	
4	6.388	
6	6.087	
1	6.687	
10	6.258	
7	6.232	
11	6.039	
3	6.046	5.005
8	6.386	5.852

- 5 The compounds can be further evaluated in a cellular chemo- and/or radiosensitization assay, an assay measuring inhibition of endogenous PARP-1 activity in cancer cell lines and eventually in an *in vivo* radiosensitization test.



CLAIMS

1. A compound of formula (I),



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

10 n is 0, 1 or 2;

X is N or CR^7 , wherein R^7 is hydrogen or taken together with R^1 may form a bivalent radical of formula $-CH=CH-CH=CH-$;

15 R^1 is C_{1-6} alkyl or thiophenyl;

R^2 is hydrogen, hydroxy, C_{1-6} alkyl, C_{3-6} alkynyl or taken together with R^3 may form $=O$;

R^3 is a radical selected from

- 20
- $-(CH_2)_s- NR^8R^9$ (a-1),
 - $-O-H$ (a-2),
 - $-O-R^{10}$ (a-3),
 - $-S-R^{11}$ (a-4), or
 - $-C\equiv N$ (a-5),

25 wherein

s is 0, 1, 2 or 3;

R^8 , R^{10} and R^{11} are each independently selected from $-CHO$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, piperidinyl, piperidinyl C_{1-6} alkyl, piperidinyl C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyl C_{1-6} alkyl, pyrrolyl C_{1-6} alkyl, aryl C_{1-6} alkylpiperidinyl, arylcarbonyl C_{1-6} alkyl, arylcarbonylpiperidinyl C_{1-6} alkyl, haloindazolylpiperidinyl C_{1-6} alkyl, or aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and

35 R^9 is hydrogen or C_{1-6} alkyl;

or R^3 is a group of formula



wherein

t is 0, 1, 2 or 3;


5 Z is a heterocyclic ring system selected from



(c-1)




(c-2)




(c-3)




(c-4)




(c-5)



(c-6)




(c-7)




(c-8)


10



(c-9)




(c-10)



(c-11)

wherein R^{12} is hydrogen, halo, C_{1-6} alkyl, aminocarbonyl, amino, hydroxy, aryl,



15

C_{1-6} alkylamino C_{1-6} alkyloxy, C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkylamino, aryl C_{1-6} alkyl, di(phenyl C_{2-6} alkenyl), piperidinyl, piperidinyl C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl C_{1-6} alkyl, aryloxy(hydroxy) C_{1-6} alkyl, haloindazolyl, aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, aryl C_{1-6} alkylamino, morpholino, C_{1-6} alkylimidazolyl, or pyridinyl C_{1-6} alkylamino;

20

R^{13} is hydrogen, piperidinyl or aryl;

R^4 , R^5 and R^6 are each independently selected from hydrogen, halo, trihalomethyl,

trihalomethoxy, C_{1-6} alkyl, C_{1-6} alkyloxy, amino, amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino,

di(C_{1-6} alkyl)amino C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl, or C_{1-6} alkyl substituted

25

with 1, 2 or 3 substituents independently selected from hydroxy, C_{1-6} alkyloxy, or amino C_{1-6} alkyloxy; or

when R⁵ and R⁶ are on adjacent positions they may taken together form a bivalent radical of formula

- $\text{-O-CH}_2\text{-O}$ (d-1),
 $\text{-O-(CH}_2\text{)}_2\text{-O-}$ (d-2),
 -CH=CH-CH=CH- (d-3), or
 $\text{-NH-C(O)-NR}^{14}\text{=CH-}$ (d-4),
 wherein R^{14} is C_{1-6} alkyl;

aryl is phenyl, phenyl substituted with halo, C₁₋₆alkyl or C₁₋₆alkyloxy;


10


with the proviso that 3-ethyl-7-[(1*H*-imidazol-1-yl)phenylmethyl]-2(1*H*)-quinoxalinone is not included.

2. A compound as claimed in claim 1 wherein
n is 0 or 1; X is N or CR⁷, wherein R⁷ is hydrogen; R¹ is C₁₋₆alkyl; R² is hydrogen;
R³ is a radical selected from (a-1) or (a-2) or is group of formula (b-1); s is 0, 1 or 2;
R⁸ is C₁₋₆alkyl or arylC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl; t is 0, 1 or 2; Z is a
heterocyclic ring system selected from (c-1), (c-2), (c-3), (c-4), (c-5) or (c-11); R¹²
is hydrogen or C₁₋₆alkyloxyC₁₋₆alkylamino; R¹³ is hydrogen; and R⁴, R⁵ and R⁶ are
each independently selected from hydrogen, halo or C₁₋₆alkyl.


3. A compound according to claim 1 and 2 wherein
n is 0 or 1; X is N; R¹ is C₁₋₆alkyl; R² is hydrogen; R³ is a radical of formula (a-1)
or is a group of formula (b-1); s is 0; R⁸ is
25 arylC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl; t is 0; Z is a heterocyclic ring system
selected from (c-1) or (c-2); R¹² is hydrogen or C₁₋₆alkyloxyC₁₋₆alkylamino; R¹³ is
hydrogen; and R⁴, R⁵ and R⁶ are each independently selected from hydrogen or
halo.

- 30 4. A compound according to claim 1, 2 and 3 selected from compound No 5, compound No 9, compound No 2 and compound No 1.





5. A compound as claimed in any of claims 1 to 4 for use as a medicine.
- 5 6. A pharmaceutical composition comprising pharmaceutically acceptable carriers and as an active ingredient a therapeutically effective amount of a compound as claimed in claim 1 to 4.
7. A process of preparing a pharmaceutical composition as claimed in claim 6 wherein
- 10 the pharmaceutically acceptable carriers and a compound as claimed in claim 1 to 4 are intimately mixed.
8. Use of a compound for the manufacture of a medicament for the treatment of a
- PARP mediated disorder, wherein said compound is a compound of formula (I)



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereo-chemically isomeric forms thereof, wherein

20 n is 0, 1 or 2;

X is N or CR⁷, wherein R⁷ is hydrogen or taken together with R¹ may form a bivalent radical of formula -CH=CH-CH=CH-;

25 R¹ is C₁₋₆alkyl or thiophenyl;

R² is hydrogen, hydroxy, C₁₋₆alkyl, C₃₋₆alkynyl or taken together with R³ may form =O;

R^3 is a radical selected from

- 5 $-(CH_2)_s-NR^8R^9$ (a-1),
 $-O-H$ (a-2),
 $-O-R^{10}$ (a-3),
 $-S-R^{11}$ (a-4), or
 $-C\equiv N$ (a-5),

wherein

s is 0, 1, 2 or 3;

- 10 R^8, R^{10} and R^{11} are each independently selected from $-CHO$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl, piperidinyll C_{1-6} alkylaminocarbonyl, piperidinyll C_{1-6} alkyl, piperidinyll C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyl C_{1-6} alkyl, 15 pyrrolyll C_{1-6} alkyl, aryll C_{1-6} alkylpiperidinyll C_{1-6} alkyl, arylcarbonyll C_{1-6} alkyl, arylcarbonylpiperidinyll C_{1-6} alkyl, haloindazolylpiperidinyll C_{1-6} alkyl, or aryll C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and R^9 is hydrogen or C_{1-6} alkyl;


or R^3 is a group of formula

- 20 $-(CH_2)_t-Z-$ (b-1),


wherein

t is 0, 1, 2 or 3;

Z is a heterocyclic ring system selected from



wherein R¹² is hydrogen, halo, C₁₋₆alkyl, aminocarbonyl, amino, hydroxy, aryl,




- 5 C₁₋₆alkylaminoC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkyloxyC₁₋₆alkylamino, arylC₁₋₆alkyl, di(phenylC₂₋₆alkenyl), piperidinyl, piperidinylC₁₋₆alkyl, C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkylC₁₋₆alkyl, aryloxy(hydroxy)C₁₋₆alkyl, haloindazolyl, arylC₁₋₆alkyl, arylC₂₋₆alkenyl, arylC₁₋₆alkylamino, morpholino, C₁₋₆alkylimidazolyl, or pyridinylC₁₋₆alkylamino;
 10 R¹³ is hydrogen, piperidinyl or aryl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, halo, trihalomethyl, trihalomethoxy, C₁₋₆alkyl, C₁₋₆alkyloxy, amino, aminoC₁₋₆alkyl, di(C₁₋₆alkyl)amino, di(C₁₋₆alkyl)aminoC₁₋₆alkyloxy or C₁₋₆alkyloxycarbonyl, or C₁₋₆alkyl substituted with 1, 2 or 3 substituents independently selected from hydroxy, C₁₋₆alkyloxy, or
 15 aminoC₁₋₆alkyloxy; or when R⁵ and R⁶ are on adjacent positions they may taken together form a bivalent radical of formula

- 20 -O-CH₂-O (d-1),
 -O-(CH₂)₂-O- (d-2),
 -CH=CH-CH=CH- (d-3), or
 -NH-C(O)-NR¹⁴=CH- (d-4),
 wherein R¹⁴ is C₁₋₆alkyl;

aryl is phenyl, phenyl substituted with halo, C₁₋₆alkyl or C₁₋₆alkyloxy.

- 25 9. Use according to claim 8 of a PARP inhibitor of formula (I) for the manufacture of a medicament for the treatment of a PARP-1 mediated disorder.
 10. Use according to claim 8 and 9 wherein the treatment involves chemosensitization.
 30 11. Use according to claims 8 and 9 wherein the treatment involves radiosensitization.
 12. A combination of a compound of formula (I) with a chemotherapeutic agent



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

5

n is 0, 1 or 2;

X is N or CR^7 , wherein R^7 is hydrogen or taken together with R^1 may form a bivalent radical of formula $-CH=CH-CH=CH-$;

10

R^1 is C_{1-6} alkyl or thiophenyl;

R^2 is hydrogen, hydroxy, C_{1-6} alkyl, C_{3-6} alkynyl or taken together with R^3 may form $=O$;

15 R^3 is a radical selected from

$-(CH_2)_s- NR^8R^9$ (a-1),

$-O-H$ (a-2),

$-O-R^{10}$ (a-3),

$-S-R^{11}$ (a-4), or

20

$-C\equiv N$ (a-5),

wherein

s is 0, 1, 2 or 3;

R^8 , R^{10} and R^{11} are each independently selected from $-CHO$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino,

25 di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl,

piperidinyl C_{1-6} alkylaminocarbonyl, piperidinyl, piperidinyl C_{1-6} alkyl,

piperidinyl C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyl C_{1-6} alkyl,

pyrrolyl C_{1-6} alkyl, aryl C_{1-6} alkylpiperidinyl, arylcarbonyl C_{1-6} alkyl,

arylcarbonylpiperidinyl C_{1-6} alkyl, haloindazolylpiperidinyl C_{1-6} alkyl, or

30 aryl C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and

R^9 is hydrogen or C_{1-6} alkyl;

or R^3 is a group of formula


$-(CH_2)_t-Z-$ (b-1),

wherein

35


t is 0, 1, 2 or 3;

Z is a heterocyclic ring system selected from



wherein R^{12} is hydrogen, halo, C_{1-6} alkyl, aminocarbonyl, amino, hydroxy, aryl,

10



15

C_{1-6} alkylamino C_{1-6} alkyloxy, C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkylamino,
 aryl C_{1-6} alkyl, di(phenyl C_{2-6} alkenyl), piperidinyl, piperidinyl C_{1-6} alkyl,
 C_{3-10} cycloalkyl, C_{3-10} cycloalkyl C_{1-6} alkyl, aryloxy(hydroxy) C_{1-6} alkyl, haloindazolyl,
 aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, aryl C_{1-6} alkylamino, morpholino, C_{1-6} alkylimidazolyl,
 or pyridinyl C_{1-6} alkylamino;
 R^{13} is hydrogen, piperidinyl or aryl;

20

R^4 , R^5 and R^6 are each independently selected from hydrogen, halo, trihalomethyl,
 trihalomethoxy, C_{1-6} alkyl, C_{1-6} alkyloxy, amino, amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino,
 di(C_{1-6} alkyl)amino C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl, or C_{1-6} alkyl substituted
 with 1, 2 or 3 substituents independently selected from hydroxy, C_{1-6} alkyloxy, or
 amino C_{1-6} alkyloxy; or
 when R^5 and R^6 are on adjacent positions they may taken together form a bivalent
 radical of formula


25



-CH=CH-CH=CH- (d-3), or
 -NH-C(O)-NR¹⁴=CH- (d-4),
 wherein R¹⁴ is C₁₋₆alkyl;

5 aryl is phenyl, phenyl substituted with halo, C₁₋₆alkyl or C₁₋₆alkyloxy.

13. A combination of a compound of formula (I) with ionizing radiation



10

the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

n is 0, 1 or 2;

15

X is N or CR⁷, wherein R⁷ is hydrogen or taken together with R¹ may form a bivalent radical of formula -CH=CH-CH=CH-;

R¹ is C₁₋₆alkyl or thiophenyl;

20

R² is hydrogen, hydroxy, C₁₋₆alkyl, C₃₋₆alkynyl or taken together with R³ may form =O;

R³ is a radical selected from

- | | | |
|----|--|-----------|
| 25 | -(CH ₂) _s -NR ⁸ R ⁹ | (a-1), |
| | -O-H | (a-2), |
| | -O-R ¹⁰ | (a-3), |
| | -S-R ¹¹ | (a-4), or |
| | -C≡N | (a-5), |

wherein

30

s is 0, 1, 2 or 3;

R⁸, R¹⁰ and R¹¹ are each independently selected from -CHO, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, amino, C₁₋₆alkylamino, di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonylaminoC₁₋₆alkyl, piperidinylC₁₋₆alkylaminocarbonyl, piperidinyl, piperidinylC₁₋₆alkyl, 35 piperidinylC₁₋₆alkylaminocarbonyl, C₁₋₆alkyloxy, thiophenylC₁₋₆alkyl,

pyrrolylC₁₋₆alkyl, arylC₁₋₆alkylpiperidinyl, arylcarbonylC₁₋₆alkyl, arylcarbonylpiperidinylC₁₋₆alkyl, haloindazolylpiperidinylC₁₋₆alkyl, or arylC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl; and R⁹ is hydrogen or C₁₋₆alkyl;

5 or R³ is a group of formula



wherein

t is 0, 1, 2 or 3;

Z is a heterocyclic ring system selected from


10




(c-1)




(c-2)




(c-3)




(c-4)




(c-5)




(c-6)




(c-7)




(c-8)



(c-9)




(c-10)



(c-11)

15

wherein R¹² is hydrogen, halo, C₁₋₆alkyl, aminocarbonyl, amino, hydroxy, aryl,



C₁₋₆alkylaminoC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkyloxyC₁₋₆alkylamino,

20

arylC₁₋₆alkyl, di(phenylC₂₋₆alkenyl), piperidinyl, piperidinylC₁₋₆alkyl,

C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkylC₁₋₆alkyl, aryloxy(hydroxy)C₁₋₆alkyl, haloindazolyl,

arylC₁₋₆alkyl, arylC₂₋₆alkenyl, arylC₁₋₆alkylamino, morpholino, C₁₋₆alkylimidazolyl,



or pyridinylC₁₋₆alkylamino;

R¹³ is hydrogen, piperidinyl or aryl;

25

R^4 , R^5 and R^6 are each independently selected from hydrogen, halo, trihalomethyl, trihalomethoxy, C_{1-6} alkyl, C_{1-6} alkyloxy, amino, amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl, or C_{1-6} alkyl substituted with 1, 2 or 3 substituents independently selected from hydroxy, C_{1-6} alkyloxy, or amino C_{1-6} alkyloxy; or


when R^5 and R^6 are on adjacent positions they may taken together form a bivalent radical of formula




wherein R^{14} is C_{1-6} alkyl;

aryl is phenyl, phenyl substituted with halo, C_{1-6} alkyl or C_{1-6} alkyloxy.


14. A process for preparing a compound as claimed in claim 1, characterized by
- a) the hydrolysis of intermediates of formula (VIII), according to art-known methods, by submitting the intermediates of formula (VIII) to appropriate reagents, such as, tinchloride, acetic acid and hydrochloric acid, in the presence of a reaction inert solvent, e.g. tetrahydrofuran.



- b) the cyclization of intermediates of formula (X), according to art-known cyclizing procedures into compounds of formula (I) wherein X is CH, herein referred to as compounds of formula (I-j), preferably in the presence of a suitable Lewis Acid, e.g. aluminum chloride either neat or in a suitable solvent such as, for example, an aromatic hydrocarbon, e.g. benzene, chlorobenzene, methylbenzene and the like; halogenated hydrocarbons, e.g. trichloromethane, tetrachloromethane and the like; an ether, e.g. tetrahydrofuran, 1,4-dioxane and the like or mixtures of such solvents.




- c) the condensation of an appropriate ortho-benzenediamine of formula (XI) with an ester of formula (XII) wherein R^h is C_{1-6} alkyl, into compounds of formula (I), wherein X is N, herein referred to as compounds of formula (I-i), in the presence of a carboxylic acid, e.g. acetic acid and the like, a mineral acid such as, for example hydrochloric acid, sulfuric acid, or a sulfonic acid such as, for example, methane-sulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid and the like.



- d) optionally effecting one or more of the following conversions in any desired order:
- (i) converting a compound of formula (I) into a different compound of formula (I);
 - (ii) converting a compound of formula (I) into the corresponding acceptable salt or N-oxide thereof;
 - (iii) converting a pharmaceutically acceptable salt or N-oxide of a compound of formula (I) into the parent compound of formula (I);
 - (iv) preparing a stereochemical isomeric form of a compound of formula (I) or a pharmaceutically acceptable salt or N-oxide thereof.

15. Use of a compound to identify PARP, wherein said compound is a compound of formula (I)



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

n is 0, 1 or 2;

5

X is *N* or CR^7 , wherein R^7 is hydrogen or taken together with R^1 may form a bivalent radical of formula $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$;

R^1 is C_{1-6} alkyl or thiophenyl;

10

R^2 is hydrogen, hydroxy, C_{1-6} alkyl, C_{3-6} alkynyl or taken together with R^3 may form $=\text{O}$;

R^3 is a radical selected from

- 15
- $-(\text{CH}_2)_s-\text{NR}^8\text{R}^9$ (a-1),
 - $-\text{O}-\text{H}$ (a-2),
 - $-\text{O}-\text{R}^{10}$ (a-3),
 - $-\text{S}-\text{R}^{11}$ (a-4), or
 - $-\text{C}\equiv\text{N}$ (a-5),

wherein

20

s is 0, 1, 2 or 3;

R^8 , R^{10} and R^{11} are each independently selected from $-\text{CHO}$, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonylamino C_{1-6} alkyl, piperidinyll C_{1-6} alkylaminocarbonyl, piperidinyll, piperidinyll C_{1-6} alkyl, piperidinyll C_{1-6} alkylaminocarbonyl, C_{1-6} alkyloxy, thiophenyll C_{1-6} alkyl, pyrrolyll C_{1-6} alkyl, aryll C_{1-6} alkylpiperidinyll, arylcarbonyll C_{1-6} alkyl, arylcarbonyllpiperidinyll C_{1-6} alkyl, haloindozolyllpiperidinyll C_{1-6} alkyl, or aryll C_{1-6} alkyl(C_{1-6} alkyl)amino C_{1-6} alkyl; and

25

R^9 is hydrogen or C_{1-6} alkyl;

30

or R^3 is a group of formula





wherein

t is 0, 1, 2 or 3;

Z is a heterocyclic ring system selected from


35





5



wherein R^{12} is hydrogen, halo, C_{1-6} alkyl, aminocarbonyl, amino, hydroxy, aryl,



- C_{1-6} alkylamino C_{1-6} alkyloxy, C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkylamino, aryl C_{1-6} alkyl, di(phenyl C_{2-6} alkenyl), piperidiny, piperidiny C_{1-6} alkyl,
- 10 C_{3-10} cycloalkyl, C_{3-10} cycloalkyl C_{1-6} alkyl, aryloxy(hydroxy) C_{1-6} alkyl, haloindazolyl, aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, aryl C_{1-6} alkylamino, morpholino, C_{1-6} alkylimidazolyl, or pyridiny C_{1-6} alkylamino;
- R^{13} is hydrogen, piperidiny or aryl;

- 15 R^4 , R^5 and R^6 are each independently selected from hydrogen, halo, trihalomethyl, trihalomethoxy, C_{1-6} alkyl, C_{1-6} alkyloxy, amino, amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl, or C_{1-6} alkyl substituted with 1, 2 or 3 substituents independently selected from hydroxy, C_{1-6} alkyloxy, or amino C_{1-6} alkyloxy; or

- 20 when R^5 and R^6 are on adjacent positions they may taken together form a bivalent radical of formula



wherein R^{14} is C_{1-6} alkyl;

aryl is phenyl, phenyl substituted with halo, C_{1-6} alkyl or C_{1-6} alkyloxy.

20. 11. 2003


(61)

ABSTRACT

5

7-PHENYLALKYL SUBSTITUTED 2-QUINOLINONES AND 2-
QUINOXALINONES AS POLY(ADP-RIBOSE) POLYMERASE INHIBITORS

- 10 The present invention provides compounds of formula (I), their use as PARP inhibitors as well as pharmaceutical compositions comprising said compounds of formula (I)



- 15 wherein n, R¹, R², R³, R⁴, R⁵, R⁶ and X have defined meanings.